Question

In the titration of sodium hydroxide and tetraoxosulphate (vi) acid, the pH at the end point is usually _____.

Answer

less than 7

Question

X is not a primary standard, but can be used to prepare a standard solution. X could be _____.

Answer

\(H_{2}C_{2}O_{4}\)

Question

Which of these substances is not employed in preparing standard solution?

Answer

Hydrochloric acid

Question

What is the concentration of a solution containing 2.5g of potassium hydroxide in (250cm^{3}) of solution? (K=39, H=1, O=16)

Answer

 $0.179 \pmod{m^{3}}$

Question

Calculate the oxidation number of sulphur in $\(FeSO_{4}.7H_{2}0\)$?

Answer

6

Question

A stock bottle of an acid reads: 56% by mass and 1.25 specific gravity. If themolar mass of the acid is 70g, find the mole per cubic decimeter.

Answer

 $10.0 \pmod{m^{3}}$

Ouestion

Which of the following is untrue of a standard solution?

Answer

Its concentration is unknown

Question The most commonly used volumetric flask in volumetric analysis is the volume capacity.
Answer 250ml
Question Pipettes are of various capacities ranging from
Answer 1ml - 50ml
Question Which of the following is not true of a desicator?
Answer it is not use for drying
Question Which of the following is not true of\((KMnO_{4}\))?
Answer it is a primary standard
Question The oxidation number of sodium in sodium hydroxide is
Answer 1
Question Oxidizing agents are
Answer electron acceptors
Question $(CO + H_{2}SO_{4})$. In the reaction above, which of the substance will be reduced?
Answer S

How many distinctive processes is involved in Iodometry?

Question

Answer two
Question The reductant in iodometry is usually
Answer thiosulphates
Question In a redox reaction, the element with less positive potential is normally
Answer reduced
Question Which of the following is least reactive?
Answer Cu
Question Neutralization reactions is usually
Answer exothermic
Question In KJ/Mol, the heat evolved when 1mole of sulphuric acid react with sodium hydroxide is approximately
Answer <57.5
Question In the titration of sodium hydroxide and tetraoxosulphate (vi) acid, the pH at the end point is usually
Answer less than 7
Question X is not a primary standard, but can be used to prepare a standard solution. X could be

```
Answer
\(H_{2}C_{2}O_{4}\)
Question
Which of these substances is not employed in preparing standard solution?
Answer
Hydrochloric acid
Question
What is the concentration of a solution containing 2.5g of potassium
hydroxide in (250 \text{cm}^{3}) of solution? (K=39, H=1, O=16)
Answer
0.179 \pmod{m^{3}}
Question
Calculate the oxidation number of sulphur in \(Feso {4}.7H {2}O\)?
Answer
Question
A stock bottle of an acid reads: 56% by mass and 1.25 specific gravity.
If themolar mass of the acid is 70g, find the mole per cubic decimeter.
Answer
10.0 \pmod{m^{3}}
Question
Which of the following is untrue of a standard solution?
Answer
Its concentration is unknown
Question
The most commonly used volumetric flask in volumetric analysis is the
_____ volume capacity.
Answer
250ml
Question
Pipettes are of various capacities ranging from _____.
```

```
1ml - 50ml
Question
Which of the following is not true of a desicator?
it is not use for drying
Question
Which of the following is not true of\(KMnO_{4}\)?
it is a primary standard
Question
The oxidation number of sodium in sodium hydroxide is _____.
Answer
Question
Oxidizing agents are .
Answer
electron acceptors
Question
\CO + H_{2}SO_{4}\). In the reaction above, which of the substance will
be reduced?
Answer
Question
How many distinctive processes is involved in Iodometry?
Answer
two
Question
The reductant in iodometry is usually _____.
Answer
thiosulphates
```

Answer

Question In a redox reaction, the element with less positive potential is normally
Answer reduced
Question Which of the following is least reactive?
Answer Cu
Question Neutralization reactions is usually
Answer exothermic
Question In KJ/Mol, the heat evolved when 1mole of sulphuric acid react with sodium hydroxide is approximately

Answer <57.5