1. Thus using starch as indicator, the colour variations of the solution depend on formation of the starch with iodide ion formed in the last stage of the titration.		
>> Complex		
Coloid		
Hydrolysed		
Navy blue		
2. The study of thermal changes in chemical and physical processes is known as		
thermochemistry		
>> Thermodynamics		
A and B		
B only		
3. In the, standard solutions of iodine are used to estimate directly the concentrations of some oxidizable species.		
>> Direct method		
indirect method		
standard method		
calibrated method		
4. The starch solution forms acomplex with the tri-iodide ion during the titration that is rapidly discharge at the end point		
Light blue		
Blue		
>> Blue black		
Brick red		
5. Theis the amount of heat released or observed for a given amount of reactants or products		
>> heat of reaction		
heat of combustion		
heat of neutralization		
heat of formation		

===== CHM191

detection,	reason and in order to avoid complications in the end point the starch indicator is added at the colouration of the ich is near the end point
	Yellow
>>	Light yellow
	Colourless
	Dark yellow
7. The most is	commonly used external indicator for iodine titration
	Glucose solution
	hydrogen chloride
	Protein solution
>>	starch solution
	he oxidizable species available for use, iodine titration are d with the $\underline{\hspace{1cm}}$
	sulphate ions
>>	Thiopsulphate
	hydrogensulphate
	thioesters
9. The oxidation potentials of Na is	
	11
	23
	271
>>	2.71
10. Complete the colour change in iodometry analysis(initially)â†'	
	Red-orange
	Dark-brown
	Brown
>>	Red-brown