===== CHM1 92		
=====		
1. The ratio of mass measured in grams per the molar mass of the substance is called		
>> mole		
concentration		
Normality		
Percentage of weight		
2. Calculate the volume of acid needed to prepare 500ml of 2.0 molar concentration of acid, if the stock bottle of the acid reads 1.25 specific gravity, 56% by mass and molar mass of acid 56.	.C	
2ml		
>> 8ml		
4ml		
10ml		
3. A solution contains 2.65g of anhydrous sodium carbonate in 250cm3 of solution. Calculate the concentration.		
10.6g/dm3		
0.1mol/dm3		
>> all of the above		
none of the above		
4. A standard solution has a known		
molar mass		
>> concentration		
molarity		
normality		
5. Gelatinous precipitate such as Aluminium hydroxide are best filtered using		
>> ashless filter paper		
sintered glass crucible		

ash filter paper

oxalate

6. The puri	ty of solid after crystallization can be verified by
	Solubility
	Ionization
>>	Taking the melting point
	Increasing the temperature
7. Impure s	olid products of chemical reactions can best be purified by
	Distillation
>>	Recrystallization
	Fractional distillation
	Sublimation
	e of the following is not a factor to be considered in good recrystallization solvent?
>>	Hardness
	Flammability
	Toxicity
	Cost
9. The most	polar solvent used for recrystallization is
>>	Water
	Ethanol
	Diethyl ether
	Methanol
	ble recrystallization solvent should possess the following cept
from the pu	It should be partially volatile in order to be easily removed rified crystals
>>	It should react with the compound being purified.
compound be	It should have boiling point below the melting point of ing purified
	It should not react with compound being purified